

Inhomogeneous phase of a Gluon Plasma at finite temperature and density

P. Castorina and D.Zappalà

*Dipartimento di Fisica, Università di Catania,
and INFN Sezione di Catania, Via Santa Sofia 64 I-95123 Catania, Italia*

(Dated: February 2, 2008)

Abstract

By considering the non perturbative effects associated with the fundamental modular region, a new phase of a Gluon Plasma at finite density is proposed. It corresponds to the transition from glueballs to non perturbative gluons which condense at a non vanishing momentum. In this respect the proposed phase is analogous to the color superconducting LOFF phase for fermionic systems.

PACS numbers: 25.75.-q, 25.75.Dw, 25.75.Nq

Quantum Chromodynamics (QCD) under extreme conditions has been intensively studied and a rich phase diagram in $T - \mu$ plane is now well established.

At small chemical potential, there is a critical temperature, T_c , where the string tension goes to zero and there is a crossover to the deconfined quark - gluon plasma (QGP). Moreover the lattice simulations of the QGP phase transition clearly indicate non perturbative effects above T_c [1, 2, 3, 4, 5, 6, 7, 8, 9, 10], that is: a) the Stefan-Boltzman limits for the pressure and the energy density of the system are not yet reached at $T \simeq 4T_c$; b) correlated $\bar{q}q$ bosonic pairs survive up to $T \simeq 2.3T_c$; c) for $T < T_c$ the gluon condensate is temperature independent whereas for $T > T_c$ its chromoelectric part rapidly decreases and, up to $2T_c$, the chromomagnetic contribution remains almost constant. On the other side of the phase diagram, i.e. at small temperature and above a critical quark chemical potential, μ_c , there is a transition to a color superconducting phase, which turns out to be stable in the so called LOFF phase, where the fermion pairs condense in a state with total momentum $\vec{q} \neq 0$ [11, 12, 13, 14, 15, 16, 17, 18].

Phenomenological quasi-particle models have been applied to fit lattice results [19, 20, 21, 22] and, in particular, for a gluon plasma (GP), i.e. a pure SU(3) gauge theory at finite temperature, lattice data on pressure and energy density can be fitted [23], for $T > 2T_c$, by a gluon gas with

the following gluon dispersion relation

$$E(k) = \sqrt{\vec{k}^2 + \frac{M^4}{\vec{k}^2}} \quad (1)$$

where $M \simeq 0.7$ GeV [23]. Equation (1) has been derived in [24, 25] after an analysis of the physical configurations of a non abelian gauge theory, once the Coulomb gauge condition, $\partial_i A_i = 0$, is fixed. This condition still leaves the freedom of having gauge equivalent configurations (or Gribov copies) which should not be counted when enumerating the physical states. The corresponding reduction to the physical states only, the so called Fundamental Modular Region (FMR), has the effect of changing the massless dispersion relation of the gluon, $E(k) = |\vec{k}|$, into Eq.(1).

The dispersion relation in Eq.(1) is extremely interesting not only for the theoretical reasons associated with confinement [26] but also because the energy has a minimum at a finite value of the momentum, $|\vec{k}|_g = M$, and increases both in the infrared and in the ultraviolet regions of the momentum. These are exactly the general conditions discussed by Brazovskii [27] which imply a stripe phase, i.e. a boson condensate in a state with $\vec{k} \neq 0$. This approach has been formulated for phase transitions in condensed matter [28], which have been experimentally tested [29].

Moreover, infrared and ultraviolet behaviors analogous to Eq.(1) are typical in non-commutative self-interacting scalar field theory [30, 31, 32] with the consequence that spontaneous symmetry breaking cannot occur for a ho-

mogeneous background but only for an inhomogeneous phase, where bosons condense at non zero momentum, analogously to the LOFF phase for fermion pairs.

Therefore, a non vanishing minimum gluon energy, as predicted from Eq.(1), opens the possibility of studying non trivial dynamical effects also at finite density. In particular, in the analysis of a pure SU(3) gauge theory at finite temperature and density, the natural question arises if the dispersion relation in Eq.(1) implies that at large density there is a gluon stripe phase. In this letter we shall address this issue.

Since from Eq.(1) the minimum energy of the gluon is $E_g^* = \sqrt{2}M$, then at $T = 0$ and $\mu = 0$ we can derive a rough estimate of glueball mass, where two valence gluons are bound by a one gluon effective exchange interaction [33]. Thus the corresponding glueball mass is

$$M_G \simeq 2E_g^* - \frac{\alpha_s(r)}{r} \quad (2)$$

where the typical scale of the bound state is $r \simeq 1/M$ and the one loop expression for $\alpha_s(r)$ has been used, with $\Lambda_{QCD} \simeq 200$ Mev. For the fitted value of M in [23], $M = 0.7$ Gev, $M_g \simeq 1.5$ Gev. This suggests that the zero momentum glueballs condensate, associated with $\langle \alpha_s/\pi G_{\mu\nu}^a G^{a\mu\nu} \rangle_0 \neq 0$, corresponds to confined gluons of energy E_g^* or minimum momentum $|\vec{k}|_g = M$ and that, by increasing the density, one can expect (see for example ref. [34]) a transition from a glueballs condensate to a deconfined, but still non perturbative, gluonic phase with condensation in the mode $|\vec{k}|_g$.

We are mainly interested in understanding the qualitative behavior of the phase diagram of the system and therefore the approximated values of the critical temperature and of the critical density can be evaluated by comparing the pressure in the two phases, and determining the transition line at $p_g - B = 0$ where

$$p_g = -TD_g \int \frac{d^3k}{(2\pi)^3} \ln[1 - e^{-\beta(E(k)-\mu)}] \quad (3)$$

is the gluon pressure, μ is the gluon (color independent) chemical potential, $D_g = 16$ is the gluon degeneracy factor and B is the bag constant that in the pure gauge theory one identifies




FIG. 1: Critical line in the $T-\mu$ plane. The minimum critical temperature corresponds to $\mu = \sqrt{2}M$.

with $\langle \alpha_s/\pi G_{\mu\nu}^a G^{a\mu\nu} \rangle$. At $\mu = 0$, the critical temperature turns out to be $T_c \simeq 0.29$ Gev for $\langle \alpha_s/\pi G^2 \rangle_0 \simeq 0.005$ Gev⁴, which is smaller than the average value obtained by QCD sum rules but still within the phenomenological uncertainty [35].

By increasing the chemical potential one obtains the critical lines in the $T - \mu$ plane, depicted in Fig.1. At a certain value of the temperature, T_c^* , the critical line reaches the maximum value of of the chemical potential allowed by Eq.(3), $\mu_c = E_g^*$. In fact for $\mu > \mu_c$ the integral in Eq.(3) is ill defined. Then, according to the standard picture of the Bose-Einstein condensation (see e.g. [36]), for $T < T_c^*$ and $\mu = \mu_c$, which corresponds to the vertical ending piece of the critical line in Fig.1, gluons progressively condense in the state of minimum energy E_g^* . When the temperature reaches the point $T = 0$, the totality of the gluons is in this state. Unlike the standard condensation where the condensed bosons carry no momentum, in this framework the gluons, after condensation, have non vanishing momentum $|\vec{k}|_g = M$.

The energy density of the system is given by

$$\epsilon_g = D_g \int \frac{d^3k}{(2\pi)^3} \frac{1}{[e^{+\beta(E(k)-\mu)} - 1]} + B \quad (4)$$




FIG. 2: Comparison of $(\epsilon_g - 3p_g)/T^4$ evaluated from Eqs.(1,2)(green points) with lattice data, for $T > 1.2T_c$. The red points correspond to the calculations with $B = 0$ [23].

and the “interaction measure” $(\epsilon_g - 3p_g)/T^4$ can be evaluated and compared with lattice data for $\mu = 0$.

The comparison is shown in Fig.2 where our results, the green points, are plotted together with those obtained in [23] and the lattice findings. Our points are evaluated for $T > 1.2T_c$, since a quasi-particle approach is unreliable near the critical point and the introduction of the gluon condensate clearly improves the agreement with respect to the case $B = 0$ [23].

The results in Fig.1 have been obtained by considering a μ independent condensate $\langle \alpha_s/\pi G^2 \rangle_0$. In fact the effect of a finite density on the value of the condensate should be taken into account. Some indications of the influence of a finite density can be obtained by following the analysis in performed in [37] where $\langle \alpha_s/\pi G^2 \rangle_\mu$ is estimated for a system with finite baryon number. As a rough evaluation we considered the same density dependence of ref. [37] including the color-flavor factor $11/3 = 11(N_c N_f)/(11N_c - 2N_f)$ with the initial value at $\mu = 0$ corresponding to the previously used value $B = \langle \alpha_s/\pi G^2 \rangle_0 \simeq 0.005 \text{ GeV}^4$. There is no qualitative difference with

respect to the previous case but since $B(\mu)$ decreases by increasing the density, the transition line is obviously characterized by a smaller critical chemical potential.

In our opinion, the previous considerations give qualitative but clear indications that there is a critical line in the $T - \mu$ plane where Eq.(1) leads to a phase transition to an inhomogeneous condensate. As firstly realized by Brazovskii, for systems in which the fluctuation spectrum has a minimum at a non zero momentum, p_c , there is a first order transition to a stripe phase, i.e. a periodic ordered state with spatial period $2\pi/p_c$. In Brazovskii’s approach the minimum in the inverse propagator at non zero momentum is determined by a self-consistent Hartree approximation and by expanding $E(k)$ around its minimum one obtains an effective lagrangian. For example, the Brazovskii-like 4-dimensional effective lagrangian for a complex scalar field can be written as [28, 30]

$$S_{eff} = \int d^4x (\alpha |\partial^2 + p_c^2| \phi|^2 + \beta |\phi|^2 + \frac{\gamma}{2} |\phi|^4) \quad (5)$$

that for $\beta < 0$ has a classical minimum at $\phi = A \exp(ipx)$ with $|A|^2 = -\beta/\gamma$ and $|p| = p_c$. In the present analysis, the minimum in $E(k)$ is due to the QCD infrared dynamics [23, 38].

On the other hand, the use of a quasiparticle approach has strong limitations. For example, since we discuss a phase transition to an inhomogeneous phase, the general expression of the pressure, $p = T \partial \ln Z / \partial V$, should be used rather than Eq.(2) which is valid for homogeneous systems. In this respect, we are practically working under the assumption of a slowly varying background and a definite answer on the existence of a stripe phase for a gluon plasma can be addressed only by lattice simulations. By the correlators method [39] a possible signal for this transition could be that, at finite density, some gauge invariant QCD correlator, C , shows an oscillating behaviour $C = a \cos(p_c x)$ corresponding to a macroscopic occupation of the modes $|\vec{k}| = \pm p_c$.

There is also the question about the surviving of this phase when quarks are taken into account. At large quark chemical potential there is the transition to a color crystalline phase with an non uniform colored condensate. Therefore if

the gluon condensation occurs at zero momentum one should obtain a LOFF fermionic phase with a superimposed gluon uniform phase. This seems unlikely due to quark-gluon interaction. From this point of view, future lattice results which support the stripe phase for pure gauge theory can give an indirect indication of the existence of the color crystalline phase at large quark chemical potential where lattice calculations are still unreliable.

In conclusion, by considering the non perturbative effects associated with the fundamental modular region, we propose a new phase of a Gluon Plasma at finite density which corresponds to the transition from glueballs to non perturbative gluons which condense at a non vanishing momentum. It will be interesting to verify if also this phenomenon is shared between field theory and condensed matter as happens for Bose-Einstein condensation and spontaneous symmetry breaking, superconductivity and chiral symmetry, LOFF phase and quarks color crystalline phase.

Acknowledgements The authors thank M. Baldo, G.Nardulli and H.Satz for useful comments.

-
- [1] G.Boyd et al., Phys. Rev. Lett. **75**(1995) 4169.
 - [2] G.Boyd et al., Nucl. Phys. **B 469**(1996) 419.
 - [3] O.Kaczmarek et al., Phys. Rev.**D 70**(2004) 074505.
 - [4] Y.Aoki et al., Phys. Lett. **B 643**(2006) 46.
 - [5] Y.Aoki et al., Nature **443**(2006) 675.
 - [6] F.Karsch and E.Ledermann, (2003) hep-lat/0305025.
 - [7] M.Asakawa, T.Hatsuda and Y. Nakahara, Nucl. Phys. **A 715**(2003) 863.
 - [8] M.Asakawa and T.Hatsuda, Nucl. Phys. **A 721**(2003) 869.
 - [9] M.D'Elia, A.Di Giacomo and E.Meggiolaro, Phys. Lett.**B 408**(1997) 315.
 - [10] M.D'Elia, A.Di Giacomo and E.Meggiolaro, Phys. Rev. **D 67**(2003)114504.
 - [11] This has been shown by using Nambu - Jona Lasinio model with color charge, see [12, 13, 14, 15, 16, 17, 18].
 - [12] K.Rajagopal and F.Wilczek hep-ph/0011333, M.G.Alford Ann. Rev. Nucl. Part. Sci. **51**(2001)131, G.Nardulli Riv. Nuovo Cimento **25 N3**(2002)1. For more recent reviews: T.Schafer hep-ph/0509068; M.Alford and K.Rajagopal hep-ph/0606157.
 - [13] R.Casalbuoni et al., Phys. Lett. **B 627**(2005) 89.
 - [14] K.Rajagopal and R.Sharma, Phys. Rev.**D 74**(2006) 094019.
 - [15] K.Rajagopal and R.Sharma, J. Phys. G **32**(2006) S483.
 - [16] M.Ciminale et al., Phys. Lett. **B 636**(2006) 317.
 - [17] K.Fukushima, Phys. Rev.**D 73**(2006) 094016.
 - [18] M.Mannarelli, K.Rajagopal and R.Sharma, hep-ph/0702021.
 - [19] P.Levai and U.W.Heinz, Phys. Rev.**C 57**(1998) 1879.
 - [20] A.Peshier and W. Cassing, Phys. Rev. Lett. **94**(2005) 172301.
 - [21] P.Castorina and M.Mannarelli, Phys. Lett. **B 336**(2007) 336.
 - [22] P.Castorina and M.mannarelli, hep-ph/0701206.
 - [23] D.Zwanziger, Phys. Rev. Lett. **94**(2005) 182301.
 - [24] V.Gribov, Nucl. Phys. **B 139**(1978)1.
 - [25] D.Zwanziger, Nucl. Phys. **B 485**(1997)185.
 - [26] J.Greensite,S.Olejnik and D.Zwanziger, hep-lat/0411032, talk at "Quark Confinement and the Hadron Spectrum VI", Villasimius, Italy, Sept. 2004.
 - [27] S.A.Brazovski, Sov. Phys. - JETP **41**(1975) 85.
 - [28] See for example, P.C. Hohenberg and J.B.Swift, Phys. Rev.**E 52**(1995) 1828.
 - [29] L.Leibler, Macromolecules **13**(1980)1602; G.H.Fredrickson and E.Helfand, J.Chem.Phys. **87** (1987)697; F.S.Bates et al. Phys. Rev. Lett. **61**(1988) 2229; F.S.Bates et al. J.Chem.Phys. **92** (1990)6225.
 - [30] S.S.Gubser and S.L.Sondhi, Nucl. Phys. **B 605**(2001) 395.
 - [31] P.Castorina and D.Zappalà, Phys. Rev.**D 68**(2003) 065008; P.Castorina,G.Riccobene and D.Zappalà, Phys. Lett. **A 337**(2005) 3463.
 - [32] J.Ambiorn and S.Catteral Phys. Lett. **B 636**(2006) 317, W.Bietenholz,F.Hofeinz and J.Nishimura Nucl. Phys. Proc. Supp **119**(2003)941, Fortsch. Phys. **51**(2003)745.
 - [33] T.H. Hansson, K. Johnson and C. Peterson, Phys. Rev. **D 26** (1982) 2069.
 - [34] A.Drago,M. Gibilisco and C.Ratti, Nucl. Phys. **A 742**(2004)165.
 - [35] M.A. Shifman, A.I. Vainshtein and V.I.Zakharov Nucl. Phys. **B 147**(1979)385, Nucl. Phys. **B 147**(1979)448; J.Reinders,H,Rubinstein and S.Yazaki , Phys. Rep. **B 127**(1985).

- [36] H.E. Haber and H.A. Weldon, Phys. Rev. Lett. **46** (1981) 1497; Phys. Rev.**D 25**(1982) 502.
- [37] M.Baldo, P.Castorina and D.Zappalà, Nucl. Phys. **A 743**(2004)3.
- [38] D.Zwanziger, hep-ph/0610021.
- [39] A. Di Giacomo, H.G. Dosch, V.I.Shevchenko and Y.A. Simonov, Phys. Rep **372**(2002)319.